

## Durham Research Online

---

### Deposited in DRO:

01 June 2015

### Version of attached file:

Published Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Mallia, C. J. and Englert, L. and Walter, G. C. and Baxendale, I. R. (2015) 'Ethyl 2-hydroxy-2-phenyl-2-(thiazol-2-yl)acetate.', Molbank., 2015 (2). M857.

### Further information on publisher's website:

<http://dx.doi.org/10.3390/M857>

### Publisher's copyright statement:

© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).

### Additional information:

---

### Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

*Short Note*

## Ethyl 2-hydroxy-2-phenyl-2-(thiazol-2-yl)acetate

Carl J. Mallia <sup>1</sup>, Lukas Englert <sup>1</sup>, Gary C. Walter <sup>2</sup> and Ian R. Baxendale <sup>1,\*</sup>

<sup>1</sup> Department of Chemistry, Durham University, South Road, Durham DH1 3LE, UK; E-Mails: c.j.mallia@durham.ac.uk (C.J.M.); englert.lukas@web.de (L.E.)

<sup>2</sup> Syngenta CP R&D Chemistry, Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY, UK; E-Mail: gary.walter@syngenta.com

\* Author to whom correspondence should be addressed; E-Mail: i.r.baxendale@durham.ac.uk; Tel.: +44-(0)-191-33-42185.

Academic Editor: Norbert Haider

Received: 7 April 2015 / Accepted: 4 May 2015 / Published: 6 May 2015

---

**Abstract:** This short note describes the synthesis of the title compound through spontaneous aerobic oxidation of ethyl 2-phenyl-2-(thiazol-2-yl)acetate. Due to the prevalence of such functional motifs in biologically active substances, we believe the oxidation encountered highlights an important degradation pathway worthy of note.

**Keywords:** thiazole; oxidation product; heterocycles

---

### Introduction

The synthesis of thiazole containing compounds has been the focus of much research due to their importance in both pharmaceuticals [1] and agrochemicals [2].

Recently, we have reported on the synthesis of 2-substituted thiazoles through a modified Gewald reaction [3]. Serendipitously, the natural air oxidation of one of the 2-substituted thiazoles led to an interesting hydroxylated thiazole which yields a glycolate moiety. This previously unreported compound is important because of its implications regarding metabolic and environmental degradation pathways for related compounds.

The air oxidation of **1** slowly gives rise to the corresponding glycolate **3** when simply left standing and open to the atmosphere; the parent compound is stable if preserved under an inert environment. The resultant glycolate **3** can be easily isolated through simple column chromatography purification.

Some related oxygenations have been previously described, however, these processes have employed either a palladium catalyst [4] or strong bases such as  $\text{Cs}_2\text{CO}_3$  [5] in the presence of oxygen.

In this short note we wish to report on the synthesis of ethyl 2-hydroxy-2-phenyl-2-(thiazol-2-yl)acetate (**3**) through air oxidation of ethyl 2-phenyl-2-(thiazol-2-yl)acetate (**1**).


## Experimental Section

In a 2–5 mL Biotage<sup>TM</sup> microwave vial, ethyl 2-cyano-2-phenylacetate (1.46 mmol, 1 equiv.) was dissolved in trifluoroethanol (6.6 mL). After 2 min stirring, 1,4-dithian-2,5-diol (0.73 mmol, 0.5 equiv.) was added and the mixture stirred for 5 min before adding triethylamine (1.61 mmol; 1.1 equiv.) followed by stirring for a further 2 min. The vial was sealed and heated under microwave irradiation for 390 min at 60 °C (conventional heating can also be used and also takes 390 min for full conversion at 60–62 °C). After cooling, the solvent was evaporated *in vacuo* and the crude residue purified using flash chromatography on silica (EtOAc/hexanes 1:4) to obtain ethyl 2-phenyl-2-(thiazol-2-yl)acetate (**1**) as a yellow oil (0.30 g, 83%). The material should be stored under an inert atmosphere.


The product **1** was left in a sealed vial at ambient temperature and after 7 days the ratio of the degradation products was analyzed by <sup>1</sup>H-NMR (71:5:24 for **1**, **2** and **3** respectively). The ratio changed when left for longer periods (>10 months ratio was 26:7:67 for **1**, **2** and **3** respectively). The oxidation products were isolated using column chromatography with a solvent mixture of EtOAc/hexanes (1:4). Four fractions were isolated; **2** ( $R_f = 0.47$ , 12 mg, 4%) [6], **1** ( $R_f = 0.26$ , 41 mg, 14%), **3** ( $R_f = 0.20$ , 181 mg, 60%) and an unknown polymeric compound ( $R_f = 0.07$ , 48 mg, 16%).

We hypothesise that the two oxidation derivatives (**2** and **3**) are generated through initial enolisation and reactive trapping of oxygen. Even though no base is present for the deprotonation, the natural enolisation is enough for the reactive trapping of oxygen, albeit rather slowly. The resultant peroxide intermediate **1b** could then potentially cyclise onto the adjacent ester moiety forming a dioxetane which, after extrusion of  $\text{CO}_2$ , would furnish product **2** (Scheme 1) [7]. Alternatively, the peroxide intermediate **1c** could undergo homolytic cleave to form the oxygen-centred radical that abstracts a hydrogen atom to form the glycolate **3** [8]. It is also possible that compound **2** is the result of ester hydrolysis (water generated in the formation of **3**), followed by decarboxylation to yield the simple 2-benzyl thiazole. Such compounds are known to oxidise to their corresponding ketones [9] or undergo a 1,2-rearrangement to form an  $\alpha$ -hydroperoxy  $\alpha$ -alkoxy ketone which would form **2** after spontaneous decomposition [10].

An alternative mechanism, not involving the initial enolisation, would be one including an initial homolytic cleavage of the C-H bond, forming a carbon centred radical **1e** which can react with oxygen to form the peroxy-radical **1f** (Scheme 2). The peroxy-radical can either react with a hydrogen atom to form **1c** as part of the formation of **3**, or form the dioxetane intermediate to yield **2**. Similar to mechanism A, there is nothing that induces the initial homolytic cleavage to initiate the reaction, however, we are convinced that considering the long reaction time needed for the transformation, small amounts of **1a** or **1e** are naturally formed due to the acidic C-H bond present in **1**.



**Scheme 1.** Putative mechanism A for the formation of **3**.



**Scheme 2.** Putative mechanism B for the formation of **3**.

*Spectroscopic Data Compound 3 (Ethyl 2-hydroxy-2-phenyl-2-(thiazol-2-yl)acetate)*

White crystalline solid;

$^1\text{H-NMR}$  (700 MHz,  $\text{CDCl}_3$ )  $\delta/\text{ppm}$  7.82 (d,  $J = 3.2$  Hz, 1H), 7.73–7.70 (m, 2H), 7.40–7.33 (m, 4H), 4.92 (s, 1H), 4.35 (q,  $J = 7.1$  Hz, 2H), 1.28 (t,  $J = 7.2$  Hz, 3H).

$^{13}\text{C-NMR}$  (176 MHz,  $\text{CDCl}_3$ )  $\delta/\text{ppm}$  172.3(C), 171.9(C), 142.9(CH), 139.6(C), 128.8(CH), 128.3(CH), 126.7(CH), 120.7(CH), 79.5(C), 63.8( $\text{CH}_2$ ), 14.1( $\text{CH}_3$ ).


IR (neat)  $\nu$  = 3457 (broad), 3118 (w), 2982 (w), 1730 (s), 1494 (w), 1449 (w), 1242 (s), 1173 (m), 1097 (m), 1066 (m), 1012 (w), 733 (m), 699 (m)  $\text{cm}^{-1}$ .

LC-MS (acetonitrile), Rt. 2.56 min,  $m/z$  = 263.9  $[\text{M}+\text{H}]^+$ . HR-MS ( $^+\text{ESI-TOF}$ ) calculated for  $\text{C}_{13}\text{H}_{14}\text{NO}_3\text{S}$  264.0694, found 264.0689 ( $\Delta$  = 1.9 ppm).

Elemental analysis: calculated for  $\text{C}_{13}\text{H}_{13}\text{NO}_3\text{S}$  C: 59.3%, H: 4.98%, N: 5.32%; measured C: 58.96%, H: 4.95%, N: 5.24%.

Melting range: 95–97 °C (*i*PrOH).

Crystal structure, CCDC-1049429 (From *i*PrOH). CCDC 1049429 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via <http://www.ccdc.cam.ac.uk/conts/retrieving.html> (or from the CCDC, 12 Union Road, Cambridge CB2 1EZ, UK; Fax: +44 1223 336033; E-mail: [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk)).



### *Phenyl(thiazol-2-yl)methanone, 2:*

Pale yellow oil;

$^1\text{H-NMR}$  (700 MHz,  $\text{CDCl}_3$ )  $\delta/\text{ppm}$  8.49–8.45 (m, 2H), 8.10 (d,  $J$  = 3.0 Hz, 1H), 7.73 (d,  $J$  = 3.0 Hz, 1H), 7.68–7.61 (m, 1H), 7.56–7.50 (m, 2H).

$^{13}\text{C-NMR}$  (176 MHz,  $\text{CDCl}_3$ )  $\delta/\text{ppm}$  184.3(C), 168.1(C), 145.0(CH), 135.4(CH), 133.8(CH), 131.2(CH), 128.6(CH), 126.4(CH).

LC-MS (acetonitrile), Rt. 2.35 min,  $m/z$  = 186.9  $[\text{M}+\text{H}]^+$ . HR-MS ( $^+\text{ESI-TOF}$ ) calculated for  $\text{C}_{10}\text{H}_8\text{NOS}$  190.0327, found 190.0323 ( $\Delta$  = 2.1 ppm).

### Acknowledgments

We would like to acknowledge the funding and support from the Royal Society (to IRB), Syngenta (to CJM) and Erasmus Programme of European Union 2013/14 (to LE), that has enabled this work to be undertaken.

## Author Contributions

Carl J. Mallia: Experimental work, writing of manuscript, literature search and synthesis planning; Lukas Englert: Experimental work; Gary C. Walter; Synthesis planning; Ian R. Baxendale: Synthesis planning and writing of manuscript.

## Conflicts of Interest

The authors declare no conflict of interest.

## References and Notes

1. Lamberth, C.; Dinges, J. *Bioactive Heterocyclic Compound Classes: Agrochemicals*; Wiley-VCH: Weinheim, Germany, 2012.
2. Baumann, M.; Baxendale, I.R.; Ley, S.V.; Nikbin, N. An overview of the key routes to the best selling 5-membered ring heterocyclic pharmaceuticals. *Beilstein J. Org. Chem.* **2011**, *7*, 442–495.
3. Mallia, C.J.; Englert, L.; Walter G.C.; Baxendale, I.R. Thiazole formation through a modified gewald reaction. *Beilstein J. Org. Chem.* **2015**, accepted.
4. Chuang, G.J.; Wang, W.; Lee, E.; Ritter, T. A dinuclear palladium catalyst for  $\alpha$ -hydroxylation of carbonyls with O<sub>2</sub>. *J. Am. Chem. Soc.* **2011**, *133*, 1760–1762.
5. Liang, Y.-F.; Jiao, N. Highly efficient C-H hydroxylation of carbonyl compounds with oxygen under mild conditions. *Angew. Chem. Int. Ed.* **2014**, *53*, 548–552.
6. <sup>1</sup>H-NMR corresponds to literature data from Riekea, R.D.; Suha, Y.-S.; Kim, S.-H. Heteroaryl manganese reagents: Direct preparation and reactivity studies. *Tetrahedron Lett.* **2005**, *46*, 5961–5964.
7. Richardson, W.H.; Hodge, V.F.; Stiggall, D.L.; Yelvington, M.B.; Montgomery F.C. 1,2-Dioxetane intermediates in the base catalyzed decomposition of  $\alpha$ -hydroperoxy ketones. *J. Am. Chem. Soc.* **1974**, *96*, 6652–6657.
8. Sakurai, H.; Kamiya, I.; Kitahara, H.; Tsunoyama, H.; Tsukuda, T. Aerobic oxygenation of benzylic ketones promoted by a gold nanocluster catalyst. *Synlett* **2009**, 245–248.
9. Santos, A.D.; Kaïm, L.E.; Grimaud, L. Metal-free aerobic oxidation of benzazole derivatives. *Org. Biomol. Chem.* **2013**, *11*, 3282–3287.
10. Sawaki, Y.; Ogata, Y. Kinetics of the base-catalyzed decomposition of  $\alpha$ -hydroperoxy ketones. *J. Am. Chem. Soc.* **1975**, *97*, 6983–6989.

© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).